



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

limb, and refraction, be considered the days will be 187 or 188. The answer for 41° N. is *good for any* other latitude north, while the problem seems to imply that an answer for 41° is different for other latitudes.

54. Proposed by S. HART WRIGHT, M. D., A. M., Ph. D., Penn Yan, N. Y.

On latitude 40° N. $=\lambda$, when the Moon's declination is $5^\circ 23'$ N. $=\delta$, and the Sun's $9^\circ 52'$ S. $=-\delta$, how long after sunset will the two horns or cusps of the Moon's crescent (recently new) set at the same moment, the crescent with its back *down* having touched the horizon first? Semi-diameters, refraction, and parallax not considered.

I. Solution by the PROPOSER.

Let B be the celestial north pole, A the zenith, AB an arc of the meridian equal the co-latitude $=c=50^\circ$, HO a portion of the horizon, SS' and MM'' portions of the diurnal arcs of the Sun and Moon, the Sun setting at S , and the Moon at M' ; BS = the polar distance of the Sun $=BS'$, and BM' the polar distance of the Moon, and AM' the zenith distance of the Moon $=90^\circ$.

Produce the vertical circle AM' to S' , S' being the place of the Sun when the Moon sets at M' . The line joining the Moon's cusps must be at right angles to the line $M'S'$ joining the centers of the Sun and Moon, and as the horizon is at right angles to $AM'S'$, the line of the cusps must lie on the horizon and set when the Moon's center sets. Put $\angle ABS = \phi$ = Sun's hour angle when it sets, and $\angle ABS' = \theta$ = Sun's hour angle when the Moon sets, and $\angle ABM' = \psi$ = Moon's hour angle when it sets.

Then we have $\cos \phi = \tan \delta' \tan \lambda$. $\therefore \phi = 81^\circ 36' 29''$, and $\cos \psi = -\tan \delta \tan \lambda$. $\therefore \psi = 94^\circ 32' 7''$. Take an auxiliary arc χ' , and $\tan \chi' = \cos \psi \cot \delta$. $\therefore \chi' = 40^\circ 0' 1''$, then $\cot A = \sin(c - \chi') \cot \chi' \operatorname{cosec} \chi'$. $\therefore A = 82^\circ 57' 55''$. Take an auxiliary angle γ' , and $\cot \gamma' = \tan A \sin \lambda$. $\therefore \gamma' = 10^\circ 52' 2''$. Then $\cos \gamma' \cot \lambda \tan -\delta' = -\cos \gamma$. $\therefore \gamma = 101^\circ 44' 43''$, and $\angle ABS' = \gamma' + \gamma = \theta = 112^\circ 36' 45''$, and $\theta - \phi = 31^\circ 0' 16'' = 2$ hours, 4 minutes, 1 second.

NOTE. The synchronous setting or rising of the cusps of a crescent Moon, is a phenomenon which must occur frequently in the tropics, and rarely or not at all beyond latitude 45° . On the 4th of July, 1897, such a moonset was very nearly accomplished, and another, almost perfect, will occur February 22, 1898, the declinations being then as given in the problem. Few persons in the northern states have ever seen the Moon set with both horns vertical.

II. Solution by G. B. M. ZERR, A. M., Ph. D., President and Professor of Mathematics, The Russell College, Lebanon, Va.

Let O be the observer, Z his zenith, HMK Moon's path, $GCSL$ Sun's path, $TEFR$ celestial equator, $AMCB$ the horizon. Let M be the position of the Moon when setting. Then, in order that the horns may set at the same time, S , M , where S is the Sun, must be on the same meridian, $ZMSN$.

$AP = \lambda = 40^\circ$. $ME = \delta = 5^\circ 23'$ N. $SF = \delta_1 = 9^\circ 52'$ S. In the triangle



